

## IXPE and multi-wavelength observations of blazar PG 1553+113 reveal an orphan optical polarization swing

RICCARDO MIDDEI,<sup>1,2</sup> MATTEO PERRI,<sup>1,2</sup> SIMONETTA PUCCETTI,<sup>1</sup> IOANNIS LIODAKIS,<sup>3</sup> LAURA DI GESU,<sup>4</sup>  
ALAN P. MARSCHER,<sup>5</sup> NICOLE RODRIGUEZ CAVERO,<sup>6</sup> FABRIZIO TAVECCHIO,<sup>7</sup> IMMACOLATA DONNARUMMA,<sup>4</sup>  
MARCO LAURENTI,<sup>8,9</sup> SVETLANA G. JORSTAD,<sup>5,10</sup> IVÁN AGUDO,<sup>11</sup> HERMAN L. MARSHALL,<sup>12</sup> LUIGI PACCIANI,<sup>13</sup>  
DAWOON E. KIM,<sup>13,14,15</sup> FRANCISCO JOSÉ ACEITUNO,<sup>11</sup> GIACOMO BONNOLI,<sup>7,11</sup> VÍCTOR CASANOVA,<sup>11</sup>  
BEATRIZ AGÍS-GONZÁLEZ,<sup>11</sup> ALFREDO SOTA,<sup>11</sup> CAROLINA CASADIO,<sup>16,17</sup> JUAN ESCUDERO,<sup>11</sup> IOANNIS MYSERLIS,<sup>18,19</sup>  
ALBRECHT SIEVERS,<sup>18</sup> POUYA M. KOUCH,<sup>3,20</sup> ELINA LINDFORS,<sup>3</sup> MARK GURWELL,<sup>21</sup> GARRETT K. KEATING,<sup>21</sup>  
RAMPRASAD RAO,<sup>21</sup> SINCHEOL KANG,<sup>22</sup> SANG-SUNG LEE,<sup>22,23</sup> SANG-HYUN KIM,<sup>22,23</sup> WHEE YEON CHEONG,<sup>22,23</sup>  
HYEON-WOO JEONG,<sup>22,23</sup> EMMANOUIL ANGELAKIS,<sup>24</sup> ALEXANDER KRAUS,<sup>19</sup> LUCIO A. ANTONELLI,<sup>25,1</sup> MATTEO BACHETTI,<sup>26</sup>  
LUCA BALDINI,<sup>27,28</sup> WAYNE H. BAUMGARTNER,<sup>29</sup> RONALDO BELLAZZINI,<sup>27</sup> STEFANO BIANCHI,<sup>30</sup> STEPHEN D. BONGIORNO,<sup>29</sup>  
RAFFAELLA BONINO,<sup>31,32</sup> ALESSANDRO BREZ,<sup>27</sup> NICCOLÒ BUCCIANTINI,<sup>33,34,35</sup> FIAMMA CAPITANO,<sup>13</sup>  
SIMONE CASTELLANO,<sup>27</sup> ELISABETTA CAVAZZUTI,<sup>36</sup> CHIEN-TING CHEN,<sup>37</sup> STEFANO CIPRINI,<sup>38,1</sup> ENRICO COSTA,<sup>13</sup>  
ALESSANDRA DE ROSA,<sup>13</sup> ETTORE DEL MONTE,<sup>13</sup> NICCOLÒ DI LALLA,<sup>39</sup> ALESSANDRO DI MARCO,<sup>13</sup>  
VICTOR DOROSHENKO,<sup>40</sup> MICHAL DOVČIAK,<sup>41</sup> STEVEN R. EHLERT,<sup>29</sup> TERUAKI ENOTO,<sup>42</sup> YURI EVANGELISTA,<sup>13</sup>  
SERGIO FABIANI,<sup>13</sup> RICCARDO FERRAZZOLI,<sup>13</sup> JAVIER A. GARCÍA,<sup>43</sup> SHUICHI GUNJI,<sup>44</sup> KIYOSHI HAYASHIDA,<sup>45,\*</sup>  
JEREMY HEYL,<sup>46</sup> WATARU IWAKIRI,<sup>47</sup> PHILIP KAARET,<sup>29</sup> VLADIMIR KARAS,<sup>41</sup> FABIAN KISLAT,<sup>48</sup> TAKAO KITAGUCHI,<sup>42</sup>  
JEFFERY J. KOLODZIEJCZAK,<sup>29</sup> HENRIC KRAWCZYNSKI,<sup>49</sup> FABIO LA MONACA,<sup>13</sup> LUCA LATRONICO,<sup>31</sup> SIMONE MALDERA,<sup>31</sup>  
ALBERTO MANFREDI,<sup>50</sup> FRÉDÉRIC MARIN,<sup>51</sup> ANDREA MARINUCCI,<sup>36</sup> FRANCESCO MASSARO,<sup>31,32</sup> GIORGIO MATT,<sup>30</sup>  
IKUYUKI MITSUISHI,<sup>52</sup> TSUNEFUMI MIZUNO,<sup>53</sup> FABIO MULERI,<sup>13</sup> MICHELA NEGRO,<sup>54,55,56</sup> CHI-YUNG NG,<sup>57</sup>  
STEPHEN L. O'DELL,<sup>29</sup> NICOLA OMODEI,<sup>39</sup> CHIARA OPPEDISANO,<sup>31</sup> ALESSANDRO PAPITTO,<sup>25</sup> GEORGE G. PAVLOV,<sup>58</sup>  
ABEL L. PEIRSON,<sup>39</sup> MELISSA PESCE-ROLLINS,<sup>27</sup> PIERRE-OLIVIER PETRUCCI,<sup>59</sup> MAURA PILIA,<sup>26</sup> ANDREA POSSENTI,<sup>26</sup>  
JURI POUTANEN,<sup>20</sup> BRIAN D. RAMSEY,<sup>29</sup> JOHN RANKIN,<sup>13</sup> AJAY RATHEESH,<sup>13</sup> OLIVER J. ROBERTS,<sup>37</sup> ROGER W. ROMANI,<sup>39</sup>  
CARMELO SGRÒ,<sup>27</sup> PATRICK SLANE,<sup>60</sup> PAOLO SOFFITTA,<sup>13</sup> GLORIA SPANDRE,<sup>27</sup> DOUGLAS A. SWARTZ,<sup>37</sup> TORU TAMAGAWA,<sup>42</sup>  
ROBERTO TAVERNA,<sup>61</sup> YUZURU TAWARA,<sup>52</sup> ALLYN F. TENNANT,<sup>29</sup> NICHOLAS E. THOMAS,<sup>29</sup> FRANCESCO TOMBESI,<sup>62,38,63</sup>  
ALESSIO TROIS,<sup>26</sup> SERGEY S. TSYGANKOV,<sup>20</sup> ROBERTO TUROLLA,<sup>61,64</sup> JACCO VINK,<sup>65</sup> MARTIN C. WEISSKOPF,<sup>29</sup>  
KINWAH WU,<sup>64</sup> FEI XIE,<sup>66,13</sup> AND SILVIA ZANE<sup>64</sup>

<sup>1</sup>Space Science Data Center, Agenzia Spaziale Italiana, Via del Politecnico snc, 00133 Roma, Italy

<sup>2</sup>INAF Osservatorio Astronomico di Roma, Via Frascati 33, 00078 Monte Porzio Catone (RM), Italy

<sup>3</sup>Finnish Centre for Astronomy with ESO, FI-20014 University of Turku, Finland

<sup>4</sup>ASI - Agenzia Spaziale Italiana, Via del Politecnico snc, 00133 Roma, Italy

<sup>5</sup>Institute for Astrophysical Research, Boston University, 725 Commonwealth Avenue, Boston, MA 02215, USA

<sup>6</sup>Physics Department, McDonnell Center for the Space Sciences, and Center for Quantum Leaps, Washington University in St. Louis, St. Louis, MO 63130, USA

<sup>7</sup>INAF Osservatorio Astronomico di Brera, Via E. Bianchi 46, 23807 Merate (LC), Italy

<sup>8</sup>INFN - Sezione di Roma "Tor Vergata", Via della Ricerca Scientifica 1, 00133 Roma, Italy

<sup>9</sup>Space Science Data Center, SSDC, ASI, Via del Politecnico snc, 00133 Roma, Italy

<sup>10</sup>Saint Petersburg State University, 7/9 Universitetskaya nab., St. Petersburg, 199034 Russia

<sup>11</sup>Instituto de Astrofísica de Andalucía, IAA-CSIC, Glorieta de la Astronomía s/n, 18008 Granada, Spain

<sup>12</sup>MIT Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA

<sup>13</sup>INAF Istituto di Astrofisica e Planetologia Spaziali, Via del Fosso del Cavaliere 100, 00133 Roma, Italy

<sup>14</sup>Dipartimento di Fisica, Università degli Studi di Roma "La Sapienza", Piazzale Aldo Moro 5, 00185 Roma, Italy

<sup>15</sup>Dipartimento di Fisica, Università degli Studi di Roma "Tor Vergata", Via della Ricerca Scientifica 1, 00133 Roma, Italy

<sup>16</sup>Institute of Astrophysics, Foundation for Research and Technology - Hellas, Voutes, 7110 Heraklion, Greece

<sup>17</sup>Department of Physics, University of Crete, 70013, Heraklion, Greece

<sup>18</sup>Institut de Radioastronomie Millimétrique, Avenida Divina Pastora, 7, Local 20, E-18012 Granada, Spain

<sup>19</sup>Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

<sup>20</sup>Department of Physics and Astronomy, 20014 University of Turku, Finland

<sup>21</sup>Center for Astrophysics — Harvard & Smithsonian, 60 Garden Street, Cambridge, MA 02138 USA

- <sup>22</sup>*Korea Astronomy and Space Science Institute, 776 Daedeok-daero, Yuseong-gu, Daejeon 34055, Korea*
- <sup>23</sup>*University of Science and Technology, Korea, 217 Gajeong-ro, Yuseong-gu, Daejeon 34113, Korea*
- <sup>24</sup>*Section of Astrophysics, Astronomy & Mechanics, Department of Physics, National and Kapodistrian University of Athens, Panepistimiopolis Zografos 15784, Greece*
- <sup>25</sup>*INAF Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monte Porzio Catone (RM), Italy*
- <sup>26</sup>*INAF Osservatorio Astronomico di Cagliari, Via della Scienza 5, 09047 Selargius (CA), Italy*
- <sup>27</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy*
- <sup>28</sup>*Dipartimento di Fisica, Università di Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy*
- <sup>29</sup>*NASA Marshall Space Flight Center, Huntsville, AL 35812, USA*
- <sup>30</sup>*Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, Via della Vasca Navale 84, 00146 Roma, Italy*
- <sup>31</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Torino, Via Pietro Giuria 1, 10125 Torino, Italy*
- <sup>32</sup>*Dipartimento di Fisica, Università degli Studi di Torino, Via Pietro Giuria 1, 10125 Torino, Italy*
- <sup>33</sup>*INAF Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125 Firenze, Italy*
- <sup>34</sup>*Dipartimento di Fisica e Astronomia, Università degli Studi di Firenze, Via Sansone 1, 50019 Sesto Fiorentino (FI), Italy*
- <sup>35</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Firenze, Via Sansone 1, 50019 Sesto Fiorentino (FI), Italy*
- <sup>36</sup>*Agenzia Spaziale Italiana, Via del Politecnico snc, 00133 Roma, Italy*
- <sup>37</sup>*Science and Technology Institute, Universities Space Research Association, Huntsville, AL 35805, USA*
- <sup>38</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Roma “Tor Vergata”, Via della Ricerca Scientifica 1, 00133 Roma, Italy*
- <sup>39</sup>*Department of Physics and Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, Stanford, California 94305, USA*
- <sup>40</sup>*Institut für Astronomie und Astrophysik, Universität Tübingen, Sand 1, 72076 Tübingen, Germany*
- <sup>41</sup>*Astronomical Institute of the Czech Academy of Sciences, Boční II 1401/1, 14100 Praha 4, Czech Republic*
- <sup>42</sup>*RIKEN Cluster for Pioneering Research, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan*
- <sup>43</sup>*California Institute of Technology, Pasadena, CA 91125, USA*
- <sup>44</sup>*Yamagata University, 1-4-12 Kojirakawa-machi, Yamagata-shi 990-8560, Japan*
- <sup>45</sup>*Osaka University, 1-1 Yamadaoka, Suita, Osaka 565-0871, Japan*
- <sup>46</sup>*University of British Columbia, Vancouver, BC V6T 1Z4, Canada*
- <sup>47</sup>*International Center for Hadron Astrophysics, Chiba University, Chiba 263-8522, Japan*
- <sup>48</sup>*Department of Physics and Astronomy and Space Science Center, University of New Hampshire, Durham, NH 03824, USA*
- <sup>49</sup>*Physics Department and McDonnell Center for the Space Sciences, Washington University in St. Louis, St. Louis, MO 63130, USA*
- <sup>50</sup>*Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, Strada Comunale Cinthia, 80126 Napoli, Italy*
- <sup>51</sup>*Université de Strasbourg, CNRS, Observatoire Astronomique de Strasbourg, UMR 7550, 67000 Strasbourg, France*
- <sup>52</sup>*Graduate School of Science, Division of Particle and Astrophysical Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan*
- <sup>53</sup>*Hiroshima Astrophysical Science Center, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan*
- <sup>54</sup>*University of Maryland, Baltimore County, Baltimore, MD 21250, USA*
- <sup>55</sup>*NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA*
- <sup>56</sup>*Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, MD 20771, USA*
- <sup>57</sup>*Department of Physics, University of Hong Kong, Pokfulam, Hong Kong*
- <sup>58</sup>*Department of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16801, USA*
- <sup>59</sup>*Université Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France*
- <sup>60</sup>*Center for Astrophysics, Harvard & Smithsonian, 60 Garden St, Cambridge, MA 02138, USA*
- <sup>61</sup>*Dipartimento di Fisica e Astronomia, Università degli Studi di Padova, Via Marzolo 8, 35131 Padova, Italy*
- <sup>62</sup>*Dipartimento di Fisica, Università degli Studi di Roma “Tor Vergata”, Via della Ricerca Scientifica 1, 00133 Roma, Italy*
- <sup>63</sup>*Department of Astronomy, University of Maryland, College Park, Maryland 20742, USA*
- <sup>64</sup>*Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, Surrey RH5 6NT, UK*
- <sup>65</sup>*Anton Pannekoek Institute for Astronomy & GRAPPA, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands*
- <sup>66</sup>*Guangxi Key Laboratory for Relativistic Astrophysics, School of Physical Science and Technology, Guangxi University, Nanning 530004, China*

## ABSTRACT

The lower energy peak of the spectral energy distribution of blazars has commonly been ascribed to synchrotron radiation from relativistic particles in the jets. Despite the consensus regarding jet emission processes, the particle acceleration mechanism is still debated. Here, we present the first X-ray polarization observations of PG 1553+113, a high-synchrotron-peak blazar observed by the Imaging X-ray Polarimetry Explorer (IXPE). We detect an X-ray polarization degree of  $(10 \pm 2)\%$

along an electric-vector position angle of  $\psi_X = 86^\circ \pm 8^\circ$ . At the same time, the radio and optical polarization degrees are lower by a factor of  $\sim 3$ . During our IXPE pointing, we observed the first orphan optical polarization swing of the IXPE era, as the optical angle of PG 1553+113 underwent a smooth monotonic rotation by about  $125^\circ$ , with a rate of  $\sim 17$  degree per day. We do not find evidence of a similar rotation in either radio or X-rays, which suggests that the X-ray and optically emitting regions are separate or, at most, partially co-spatial. Our spectro-polarimetric results provide further evidence that the steady-state X-ray emission in blazars originates in a shock-accelerated and energy-stratified electron population.

*Keywords:* acceleration of particles, black hole physics, polarization, radiation mechanisms: non-thermal, galaxies: active, galaxies: jets

## 1. INTRODUCTION

Blazars, a rare class of active galactic nuclei (AGN; Antonucci 1993; Urry & Padovani 1995; Padovani et al. 2017), display highly time-variable radiation across the entire electromagnetic spectrum, from radio to very high-energy (TeV)  $\gamma$ -rays (e.g., Angel & Stockman 1980; Padovani & Giommi 1995; Ghisellini et al. 1998a,b; Sreekumar et al. 1998; Fossati et al. 1998; Abdo et al. 2010; Maselli et al. 2010; Acero et al. 2015; Agudo et al. 2011a,b; Liodakis et al. 2018b, 2019; Hovatta & Lindfors 2019). It is widely accepted that the nonthermal radiation originates in a highly collimated jet of relativistic particles extended along the polar axis of an accreting supermassive black hole (SMBH) and pointing toward the Earth (see e.g. Blandford et al. 2019). Despite accretion of matter onto the SMBH powering blazars, their orientation is such that highly relativistically boosted radiation from the jet dominates the emission spectrum (e.g., Lähteenmäki & Valtaoja 1999; Hovatta et al. 2009; Liodakis et al. 2018a), with a double-peaked spectral energy distribution (SED). The first hump peaks between infrared (IR) and X-ray frequencies, and is ascribed to synchrotron radiation from energetic electrons, perhaps, as suggested by previous IXPE pointings, accelerated in shocks (e.g., Liodakis et al. 2022; Di Gesu et al. 2022). The second hump is located in the GeV – TeV range, with the mechanism behind the emission still debated. Both hadronic and leptonic frameworks have been proposed (e.g. Maraschi et al. 1992; Boettcher 2012; Cerruti et al. 2015), although recent X-ray spectro-polarimetric observations seem to favor a leptonic scenario in which synchrotron self-Compton emission plays a key role in shaping the SED of some blazars (see e.g. Middei et al. 2023; Peirson et al. 2023). Depending on the frequency at which the synchrotron component peaks ( $\nu_{peak}$ ), blazars are clas-

sified as high-synchrotron-peak (HSP), intermediate-synchrotron-peak (ISP) or low-synchrotron-peak (LSP), with  $\nu_{peak} > 10^{15}$  Hz,  $10^{14} < \nu_{peak} < 10^{15}$  Hz, and  $\nu_{peak} < 10^{14}$  Hz, respectively (e.g., Ajello et al. 2020).

PG 1553+113 (RA=15h 55m 43.0440s, Dec= +11° 11' 24.365", J2000) is an HSP source ( $\nu_{peak} \approx 3.9 \times 10^{15}$  Hz, Ajello et al. 2020), therefore X-rays sample the falling part of the low-energy hump of its SED. Although several redshift values ( $z < 0.5$ ) have been reported for this BL Lacertae object, the most commonly adopted value is  $z=0.433$  (see, e.g., Nicastro et al. 2018; Johnson et al. 2019; Dorigo Jones et al. 2022). It is a TeV  $\gamma$ -ray emitter (Aleksić et al. 2012), and has exhibited one of the most compelling cases of quasi-periodic oscillations from a blazar at  $\gamma$ -ray energies as a period  $p=2.18 \pm 0.08$  (years) was found in the  $\gamma$ -rays (Ackermann et al. 2015; Tavani et al. 2018).

Here we present the first X-ray polarization observation of PG 1553+113, an HSP blazar observed by the Imaging X-ray Polarimetry Explorer (IXPE; Weisskopf et al. 2022) in 2023 February. Owing to the three gas pixel detectors (GPDs; Costa et al. 2001; Bellazzini et al. 2003; Baldini et al. 2021) in its focal plane, IXPE provides an unprecedented opportunity to measure the X-ray polarization of cosmic sources. Since its launch on 2021 December 9, IXPE has observed different types of AGNs (e.g., Marinucci et al. 2022; Ursini et al. 2023; Gianolli et al. 2023; Ingram et al. 2023; Tagliacozzo et al. 2023; Ehlert et al. 2022), including blazars (e.g. Liodakis et al. 2022; Di Gesu et al. 2022; Middei et al. 2023). PG 1553+113 is one of several HSP blazars observed by IXPE. In §2 we report on the data reduction, while §3 discusses the X-ray emission and polarization properties of PG 1553+113. In §4 we present the contemporaneous radio and optical polarization observations, and in §5 we summarize and discuss our results and draw conclusions.

\* Deceased

## 2. X-RAY DATA REDUCTION

**Table 1.** Log of the IXPE and XMM-Newton observations of PG 1553+113.

Observatory	Obs. ID	Obs. date	Net exp.
		yyyy-mm-dd/dd	ks
IXPE(a)	02004999	2023-02-01/03	~ 98
IXPE(b)	02004999	2023-02-08/09	~ 28
XMM-Newton	0902112101	2023-01-31	~14

PG 1553+113 was observed with IXPE’s three Detector Units (DUs) in 2023 February for a total exposure time of  $\sim 130$  ks. The observing time was split into two intervals (hereafter, intervals a and b). Interval a was taken in 2023 February 1-2, for a total net exposure time of  $\sim 100$  ks. Interval b, performed one week later in February 8-9, had a net exposure time of  $\sim 30$  ks.

We coordinated the IXPE observations with XMM-Newton (Jansen et al. 2001) and the Neil Gehrels Swift observatory (Swift; Gehrels et al. 2004), which observed the target before and after the IXPE observation. Table 1 reports the log for the IXPE and XMM-Newton pointings.

We extracted the  $I$ ,  $Q$  and  $U$  spectra for each of the three DUs of IXPE using the software IXPEOBSSIM (v. 30.0.0; Pesce-Rollins et al. 2019; Baldini et al. 2022) and following the background rejection prescriptions presented in Di Marco et al. (2023). Moreover, spectra were computed in order to use the so-called weighted analysis method presented in Di Marco et al. (2022, parameter STOKES = NEFF in XSELECT). The spectra for the Stokes parameters were computed from a circular region with radius= $0.95'$  centered on the source, while an annulus with  $r_{\text{in(out)}}=1.2'(3.5')$  was adopted to extract the background. This approach has been shown to enhance the sensitivity to polarization (Di Marco et al. 2023). The resulting  $I$  Stokes spectra were grouped by requiring each energy bin to have a signal-to-noise ratio greater than 7. We adopted uniform binning of 280 eV for the  $Q$  and  $U$  Stokes spectra.

XMM-Newton performed a snapshot ( $\sim 14$  ks long) of PG 1553+113, and we analyzed the data taken with the EPIC-pn camera (Strüder et al. 2001). The source was observed in Small Window mode, with the medium filter applied. We processed the data via the standard XMM-Newton Science Analysis System (SAS v21, Gabriel et al. 2004). Source extraction radii and screening for high-background intervals were determined through an iterative process (Piconcelli et al. 2004) that maximizes the signal-to-noise ratio. The background was extracted from circular regions with a radius of  $50''$ , and the same shape centered on the source PG 1553+113 was adopted

for science products. The resulting third-level products were grouped by requiring each bin to contain at least 30 counts, and not to over-sample the spectral resolution by a factor larger than 3. The net count rate was less than the maximum allowed limit of  $50 \text{ cts s}^{-1}$  to avoid deteriorated response due to photon pile-up for EPIC-pn observations in Small Window mode (e.g. Jethwa et al. 2015). We further assessed the potential impact of pile-up in the XMM-Newton observation by means of the *epatplot* task, a standard SAS command devoted to checking for any pile-up affecting the data, and we found it to be negligible.

Swift pointed at PG 1553+113 before and after the IXPE pointing. Sixteen exposures, each 1 ks long, were taken in photon counting mode and calibrated and reduced using the *XRTDAS* software package<sup>1</sup>. To extract the source spectra, we adopted a circular region of radius  $47''$ , while the background was derived with a concentric annulus with inner(outer) radii of  $120''(150'')$ . Spectra were then binned, with at least 25 counts in each bin, in order to meaningfully use the  $\chi^2$  statistic in our spectral analysis.

### 3. X-RAY ANALYSIS

The X-ray spectro-polarimetric analysis was performed using XSPEC (Arnaud 1996) and following the prescriptions of the so-called weighted analysis method (Di Marco et al. 2022). We accounted for Galactic absorption along the line of sight using the *tbabs* model. In all the fits, we set the Galactic column density to  $n_{\text{H}}=3.62 \times 10^{20} \text{ cm}^{-2}$  (HI4PI Collaboration et al. 2016). For calculations involved the distance to PG 1553+113, we adopt the standard cosmological framework with  $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$  and  $\Lambda_0 = 0.73$ .

#### 3.1. IXPE and XMM-Newton spectro-polarimetric analysis

To determine the spectral properties of PG 1553+113, we modeled the 0.3–10 keV XMM-Newton spectrum (taken contemporaneously with the IXPE observation; see Table 1). We tested two models: an absorbed power law and an absorbed logarithmic parabola. The log-parabola (Massaro et al. 2004) is defined by  $N(E) = (E/E_1)^{-\alpha - \beta \log(E/E_1)}$ , where  $\alpha$  accounts for the source photon index at energy  $E_1$ ,  $\beta$  represents the curvature of the parabola, and  $E_1$  is a reference (“pivot”) energy. In the fits, we determined the photon index and normalization for the power law. For the log-parabola, we fixed  $E_1$  to 3 keV and left  $\alpha$ ,  $\beta$ , and the normalization free to vary. The power law fails

<sup>1</sup> <https://sda2006.ts.infn.it/presentazioni/capalbi.pdf>

to reproduce the XMM-Newton spectrum satisfactorily ( $\chi^2/\text{d.o.f.}=530/159$ ), while the logarithmic parabola returns a better statistic, with  $\chi^2=219$  for 159 degrees of freedom. We obtained  $\alpha = 2.50 \pm 0.01$  and  $\beta = 0.13 \pm 0.01$ , which is compatible with a soft and curved spectrum, as expected for a blazar X-ray spectrum dominated by synchrotron radiation. We note that this fit can be further improved ( $\Delta\chi^2=-41$  for two additional degrees of freedom) by adding a narrow Gaussian component to account for an additional soft X-ray feature ( $E = 0.68$  keV and  $\text{Norm}=2.58 \times 10^{-4}$  photons  $\text{cm}^{-2} \text{s}^{-1}$ ). Finally, we notice that IXPE caught PG 1553+113 in a fairly high flux state compared with the average flux of this source ( $F_{\text{mean}} \sim 2.1 \times 10^{-11}$  erg  $\text{cm}^{-2} \text{s}^{-1}$  in the 2–10 keV energy range) obtained by [Giommi et al. \(2021\)](#) who analysed 802 Swift archival snapshots of the blazar.

Next, we included in the analysis the IXPE spectra extracted from both the a and b intervals. We assigned each spectrum to a “datagroup” in XSPEC, which allowed us to couple/decouple the parameters of the model between the different instruments when needed. As a first test, we assumed the polarization properties of PG 1553+113 to be the same in the two observing intervals. Thus, we fit the corresponding  $I$ ,  $Q$ , and  $U$  Stokes spectra while constraining the relevant parameters. To account for polarization, we updated our XSPEC model to `tbabs×const×polconst×logpar`. The constant model (`const`) allows for cross-calibration among the different IXPE DUs and the EPIC-pn camera. The polarization model `polconst`, which assumes constant polarization parameters within the operating energy range, has two free parameters: the polarization degree ( $\Pi_X$ ) and the electric-vector position angle ( $\Psi_X$ , measured from north through east). Finally, for the `logpar` model, we coupled the  $\alpha$  and  $\beta$  parameters between the IXPE and XMM-Newton spectra. This model provided a statistically acceptable ( $\chi^2 / \text{d.o.f.}=471/452$ ) fit to the IXPE and XMM-Newton data, which is shown in Fig. 1 (panels a and b for the  $I$  and  $Q$  and  $U$  Stokes parameters, respectively). The corresponding best-fit parameters are listed in Table 2. Our fit result suggests that the spectrum of PG 1553+113 had a soft and curved shape ( $\alpha = 2.49 \pm 0.01$  and  $\beta = 0.11 \pm 0.01$ ). These values are well within the range of what is commonly observed in HSP blazars (e.g. [Middei et al. 2022](#)). The polarization properties were determined to be  $\Pi_X=(10.1 \pm 2.3)\%$  and  $\psi_X = 86^\circ \pm 8^\circ$ . In Fig. 1c we show the contours corresponding to 68%, 90%, and 99% confidence levels. We find the polarized signal to be maximum in terms of significance in the 2–5.5 keV energy range. The soft and curved shape of the spectrum, and the maximum polar-


**Table 2.** Best-fit parameters from the spectro-polarimetric analysis of PG 1553+113. Time-average results refer to the IXPE and XMM-Newton joint fit, while values quoted for intervals a and b were obtained only with the IXPE data. Fluxes are in units of  $10^{-11}$  erg  $\text{cm}^{-2} \text{s}^{-1}$  while the logarithmic parabola has a normalization of  $10^{-3}$ . All errors are given at 68% confidence interval for 1 parameter of interest (i.e.  $\Delta\chi^2=1$ ), while the upper limit to the polarization degree corresponds to 99% uncertainty. The symbol † indicates those parameters that were kept frozen during the model fits.

Model	Parameter	time average	Interval a	Interval b
polconst	$\Pi_X$	$(10.1 \pm 2.3)\%$	$(11.6 \pm 3.4)\%$	$< 26\%$
	$\psi_X$	$86^\circ \pm 8^\circ$	$85^\circ \pm 8^\circ$	-
tbabs	$n_H$	3.62†	3.62†	3.62†
log-par	$\alpha$	$2.491 \pm 0.007$	$2.59 \pm 0.03$	$2.56 \pm 0.15$
	$\beta$	$0.11 \pm 0.01$	0.11†	0.11†
	Norm	$1.49 \pm 0.02$	$1.41 \pm 0.03$	$1.43 \pm 0.02$
const	$K_{\text{Du}2}$	$0.91 \pm 0.01$	$0.95 \pm 0.01$	$0.96 \pm 0.07$
const	$K_{\text{Du}3}$	$0.95 \pm 0.01$	$0.90 \pm 0.01$	$0.97 \pm 0.07$
const	$K_{\text{Epic-pn}}$	$1.04 \pm 0.01$	-	-
$F_{2-8 \text{ keV}}$		$2.55 \pm 0.03$	$2.47 \pm 0.03$	$2.44 \pm 0.05$

ization signal measured below 5.5 keV, are in agreement with the PG 1553+113 X-ray spectrum being dominated by synchrotron emission.

We further searched for a possible change in the polarization properties of PG 1553+113 between observation intervals a and b. Using the model `tbabs×const×polconst×logpar`, we computed  $\alpha$ , the normalization,  $\Pi_X$ , and  $\psi_X$  from the Stokes parameters derived for intervals a and b separately. In this test the curvature parameter  $\beta$  was kept frozen to its best-fit value indicated in Table 2<sup>2</sup>. The adoption of the `tbabs×const×polconst×logpar` model led us to a very good representation of the two data sets, with  $\chi^2/\text{d.o.f.}=304/276$  and  $\chi^2/\text{d.o.f.}=285/276$  for intervals a and b, respectively. The inferred best fit parameters are listed in Table 2. Despite the compatible 2–8 keV flux observed in the two intervals,  $\Pi_X$  and  $\psi_X$  are only constrained within interval a, while an upper limit  $\Pi_X^{99\%} < 26\%$  is obtained when fitting the Stokes  $I$ ,  $Q$  and  $U$  spectra from interval b. This upper limit is likely the result of the relatively short duration of the b interval exposure. As a final test, we studied the polarization properties of PG 1553+113 over time spans shorter than the duration of intervals a and b of the IXPE observa-

<sup>2</sup> For this test we did not consider the XMM-Newton data, as they were taken just before the IXPE interval a exposure



**Figure 1.** Top Panels: Best fit to the IXPE and XMM-Newton data and corresponding residuals. On the left, we show the model  $\text{const} \times \text{polconst} \times \text{logpar}$  fitting the  $I$  Stokes spectra, while on the right the best-fit  $Q$  and  $U$  Stokes spectra are shown. Bottom panels: regions accounting for 68%, 90%, and 99% confidence levels computed using the full IXPE 2-8 keV data (bottom left) or taking into account the IXPE  $I$ ,  $Q$  and  $U$  Stokes spectra only in the 2-5.5 keV range.

tion. However, for all the tested temporal intervals no hints of variability were found.

### 3.2. Swift monitoring campaign


A Swift campaign was organized to monitor the flux and spectral variability of PG 1553+113. The *XRT* observed the blazar from 2023 early January until late February. We fit the resulting 16 spectra (exposures of  $\leq 1$  ks each) with a log-parabola model, with Galactic absorption (see above). The adopted model provides a statistically acceptable representation of the data. We show the spectral parameters and flux estimates inferred from the fits in Fig. 2. The Swift data are compatible with nearly constant spectral shape and variability of flux by less than a factor of 2. The average values for the quantities displayed in Fig. 2 are:  $\alpha_{\text{avg}} = 2.15 \pm 0.12$  and  $\beta_{\text{avg}} = 0.3 \pm 0.2$ , with fluxes  $F_{\text{IXPE}}^{\text{avg}} = 3.4 \pm 0.6$  (2-8 keV),  $F_{\text{Soft}}^{\text{avg}} = 4.6 \pm 0.5$  (0.5-2 keV), and  $F_{\text{Hard}}^{\text{avg}} = 3.7 \pm 0.6$  (2-10 keV) in units of  $10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$ .

## 4. MULTI-WAVELENGTH OBSERVATIONS

PG 1553+113 was also observed at radio and optical wavelengths by the Effelsberg 100-m telescope, the Korean VLBI Network (KVN), Institut de Radioastronomie Millimétrique (IRAM 30-m telescope), Submillimeter Array (SMA), Boston University Perkins Telescope, Calar Alto Observatory, Nordic Optical Telescope and Sierra Nevada Observatory. The observational and data analysis procedures are described in detail in [Middei et al. \(2023\)](#). All the multiwavelength observations are available upon request to the individual observatories. We supplement our campaign with 3-day-binned publicly available Fermi Large Area Telescope data from the light curve repository<sup>3</sup> ([Abdollahi et al. 2023](#)).

At radio wavelengths, the Effelsberg observation on 2023 February 8 (MJD 59983.14) was performed at 4.85 GHz as part of the QUIVER (Monitoring the Stokes  $Q$ ,  $U$ ,  $I$  and  $V$  Emission of AGN jets in Radio) program

<sup>3</sup> <https://fermi.gsfc.nasa.gov/ssc/data/access/lat/LightCurveRepository/>




**Figure 2.** Results of Swift-XRT monitoring campaign of PG 1553+113. Fluxes  $F_{\text{IXPE}}$ ,  $F_{\text{Soft}}$ , and  $F_{\text{Hard}}$  are computed in the 2-8, 0.5-2, and 2-10 keV bands, respectively, and are displayed in units of  $10^{-11}$  erg  $\text{cm}^{-2}$   $\text{s}^{-1}$ . “HR” is the hardness ratio, defined as  $(F_{\text{Hard}} - F_{\text{Soft}}) / (F_{\text{Hard}} + F_{\text{Soft}})$ . Gray shaded areas mark the duration of the two IXPE observations, intervals a (left) and b (right). Black data points account for the different quantities measured by IXPE and quoted in Table 2.

(Kraus et al. 2003; Myserlis et al. 2018). The IRAM-30m observations were obtained and analyzed in the framework of the POLAMI (Polarimetric Monitoring of AGN at Millimeter Wavelengths) project<sup>4</sup> (Agudo et al. 2018a,b; Thum et al. 2018). The Korean VLBI Network (KVN) observation was conducted on 2023 February 8 (UT 20) with two 21-m antennas (KVN Yonsei and Tamna) combined in single-dish mode. The polarization observations were conducted in position-switching mode (Kang et al. 2015) and calibrated using an unpolarized (Jupiter) calibrator and a polarized (3C286, Agudo et al. 2012) calibrator for the polarization degree, and the Crab nebula for the polarization angle ( $152^\circ$ ; Aumont et al. 2010). The SMA observations were performed within the framework of the SMA Monitoring of AGNs with Polarization (SMAPOL) program (Ho et al. 2004; Marrone & Rao 2008; Primiani et al. 2016). The radio observations covered a frequency range from 4.85 to 225.5 GHz. At the lowest radio frequency (4.85 GHz), the polarization de-

gree was  $\Pi_{4.85\text{GHz}} = 2.6 \pm 0.7\%$  along a polarization angle  $\psi_{4.85\text{GHz}} = 133 \pm 7^\circ$ . At intermediate frequencies (22 GHz, 43 GHz), the polarization degree was  $\Pi_{22-43\text{GHz}} = 1 - 2\%$ , with  $\psi_{22-43\text{GHz}} = 65 - 80^\circ$ . At the higher frequencies (86 GHz, 225 GHz), we measured  $\Pi_{86-225\text{GHz}} \sim 3\%$  along  $\psi_{86-225\text{GHz}} = 100 - 120^\circ$ . We did not observe strong variability in either the degree or angle of polarization at any of the radio frequencies before, during, or after the IXPE observations (Fig. 3).

Optical observations were obtained at the Calar Alto Observatory, Nordic Optical Telescope, Perkins Telescope, and Sierra Nevada Observatory. At Calar Alto Observatory, we used the 2.2 m telescope and CAFOS (Calar Alto Faint Object Spectrograph). Observations at Sierra Nevada Observatory used the T90 telescope. The data for both observatories were taken in *R*-band and analyzed following standard polarimetric procedures. *B*, *V*, *R*, and *I* band polarimetric observations were obtained with the Alhambra Faint Object Spectrograph and Camera (ALFOSC) at the Nordic Optical Telescope, and analyzed using Tuorla Observatory’s semi-automatic data reduction pipeline (Hovatta et al. 2016; Nilsson et al. 2018). Additional photo-

<sup>4</sup> <http://polami.iaa.es/>



**Figure 3.** Multi-wavelength radio and optical observations of PG 1553+113. Panel a shows the brightness in magnitudes, panel b the polarization degree, panel c the polarization angle, and panel d the  $\gamma$ -ray light curve in the 0.1-100 GeV range. The gray shaded areas mark the two intervals of IXPE observations. The error bars represent  $1\sigma$  uncertainties. Pink boxes identify the  $\psi_X$  best-fit value and its  $1\sigma$  uncertainty as derived using IXPE. Panel d displays the photon flux observed by the FERMI Large Area Telescope during the IXPE monitoring of the source. The photon flux is in units of  $10^{-7}$  photons  $s^{-1} \text{ cm}^{-2}$  (see Abdollahi et al. 2023, for details of the Fermi light curve).



metric ( $B$ ,  $V$ ,  $R$ ,  $I$ ) and polarimetric ( $R$ -band) data were obtained at the Perkins Telescope (1.8 m) using the PRISM camera. During IXPE interval a, we found  $\Pi_O = 2.2 \pm 0.4\%$ . During interval b, we observed an increase of the polarization degree to  $\Pi_O = 4.2 \pm 0.5\%$ . At the same time, we observed a rotation of the polarization angle of about  $\Delta\psi_R \approx 125^\circ$  at a rate of about  $17^\circ \text{ day}^{-1}$  (see Fig. 3, panel c). The duration, amplitude and rotation rate are within typical rotation parameters of other blazars (e.g., Blinov et al. 2018). Compared to the two previous rotations of the optical polarization vector of PG 1553+113 detected by the RoboPol program (Blinov et al. 2015, 2016a, 2018), the event reported here has a similar amplitude, but a factor of 2-3 higher rotation rate. However, we note that, while the end of the rotation is apparent in our data, the rotation may have started before the beginning of our observations. It is therefore possible that we are underestimating the total amplitude and duration of the event.

## 5. DISCUSSION AND CONCLUSIONS

We have reported the first IXPE observations of the HSP blazar PG 1553+113, finding an X-ray polarization degree  $\Pi_X = (10.1 \pm 2.3)\%$  along an electric-vector position angle of  $\psi_X = 86^\circ \pm 8^\circ$ . These values were obtained by performing a spectro-polarimetric analysis, using XSPEC, of the  $I$ ,  $Q$ , and  $U$  Stokes spectra that were fit with a quasi-simultaneous XMM-Newton spectrum. The source spectrum can be fit with a log-parabola, which models the curved X-ray spectrum typical of synchrotron-dominated HSP blazars (e.g. Massaro et al. 2004; Giommi et al. 2021; Middei et al. 2022).

Our contemporaneous radio and optical polarization observations find a 2.5–5 times lower polarization degree, with a polarization angle roughly aligned with the X-ray value. Previous IXPE results for the HSP blazars Mrk 421 and Mrk 501 (Liodakis et al. 2022; Di Gesu et al. 2022) have found similar chromatic behavior of  $\Pi$ , usually with similar  $\psi$  across different frequencies. The ratio  $\Pi_X/\Pi_O$  is also similar to that found in Mrk 421 and Mrk 501. These results have been interpreted as evidence of an electron population that is accelerated in a shock and becomes energy stratified as it cools while propagating away from the shock front (see details in, e.g., Marscher & Gear 1985; Perlman et al. 1999; Perlman & Wilson 2005; Angelakis et al. 2016; Tavecchio et al. 2018). Given the different synchrotron cooling lengths, the optical and radio emitting particles occupy an increasing volume of the jet with decreasing frequency. In this case, the radio, optical, and X-ray emission regions are expected to be at most partially

co-spatial, which leads to different polarization properties in different bands.

Rotations with time of the optical polarization angle have now been observed in a large number of blazars (e.g., Marscher et al. 2008, 2010; Blinov et al. 2016b, 2018; Liodakis et al. 2020). Recently, rotation of the X-ray polarization vector was observed for the first time, in Mrk 421 (Di Gesu et al. 2023). During this event, Mrk 421 was observed to have a variable X-ray spectral shape and flux, but no evidence was found for a change in  $\psi$  at either radio or optical wavelengths. The X-ray rotation was interpreted as a shock propagating along a helical magnetic field.

In the case of PG 1553+113, we instead have observed  $\Delta\psi \approx 125^\circ$  at optical wavelengths, which occurred between IXPE intervals a and b. This rotation of the optical polarization angle was accompanied by a modestly variable X-ray flux and roughly constant spectral shape. Moreover, we have found no evidence for a change in  $\psi$  in either the radio or X-ray band. This is consistent with the interpretation of the Mrk 421 results, where the X-ray-optical–radio emission of blazars originates not from a single localized region, but rather from separate (or at most partially co-spatial) regions in the jet. Moreover, as imaged with the MOJAVE program<sup>5</sup>, PG1553+113 is a very core-dominated source at 15 GHz, with faint extended structure at a projected position angle between  $30\text{--}50^\circ$ . This implies that the observed X-ray polarization position angle is oblique to the parsec-scale jet direction, with a difference  $\sim 45^\circ$ , similar to that observed for Mrk 421 during its first IXPE observation in 2022 May (Di Gesu et al. 2023). However, we note that the position angle of the jet is known to vary over time (Lico et al. 2020).

The origin of the optical rotation is unclear. Figure 3 (bottom panel) shows the 3-day-binned  $\gamma$ -ray light curve of PG 1553+113 during the IXPE observations. There is a clear brightening of the  $\gamma$ -rays near the center of the optical polarization angle rotation. A statistical association between optical rotations and  $\gamma$ -ray activity has already been established by the RoboPol program (Blinov et al. 2015, 2018), suggesting that such rotations are often deterministic rather than a random walk of the polarization angle. However, the statistics do not exclude the possibility that some rotations are indeed random walks. Kink-driven magnetic reconnection (Bodo et al. 2021) can cause variations of the polarization angle that would be different in optical and X-rays. However, in this scenario, the optically emitting particles do

<sup>5</sup> <https://www.cv.nrao.edu/MOJAVE/sourcepages/1553+113.shtml>

not travel far from the current sheet, therefore we expect a similar polarization degree in optical and X-rays (Bodo et al. 2021). The observed chromatic behavior of  $\Pi$  disfavors this interpretation. Merging of reconnection plasmoids can also produce orphan optical polarization rotations accompanied by multiwavelength flares (Hosking & Sironi 2020). However, the timescales of such rotations are expected to be much shorter than the  $\sim 7$  days observed here. In addition, we did not observe the expected optical flare, which makes this interpretation unlikely.

We conclude that the lower degree of polarization at optical wavelengths is most likely connected with stronger turbulence in the optical emission region. In this case, the detected rotation could have been caused by turbulence, as observed in simulated polarization angle behavior in the Turbulent Extreme Multi-Zone (TEMZ) model and Particle-in-Cell simulations (Marscher et al. 2017; Marscher & Jorstad 2021; Zhang et al. 2023).

Given the above assessment, our results favor a model where an energy stratified — and most likely shock-accelerated — electron population is responsible for the non-thermal X-ray–optical–radio emission of at least some, and possibly all, astrophysical jets associated with SMBHs.

**Acknowledgments:** The Imaging X-ray Polarimetry Explorer (IXPE) is a joint US and Italian mission. The US contribution is supported by the National Aeronautics and Space Administration (NASA) and led and managed by its Marshall Space Flight Center (MSFC), with industry partner Ball Aerospace (contract NNM15AA18C). The Italian contribution is supported by the Italian Space Agency (Agenzia Spaziale Italiana, ASI) through contract ASI-OHBI-2017-12-I.0, with agreements ASI-INAF-2022-14-HH.0 and ASI-INFN 2021-43-HH.0, and its Space Science Data Center (SSDC), and by the Istituto Nazionale di Astrofisica (INAF) and the Istituto Nazionale di Fisica Nucleare (INFN) in Italy. This research used data products provided by the IXPE Team (MSFC, SSDC, INAF, and INFN) and distributed with additional software tools by the High-Energy Astrophysics Science Archive Research Center (HEASARC), at NASA Goddard Space Flight Center (GSFC). We acknowledge financial support from ASI-INAF agreement n. 2022-14-HH.0. The research at Boston University was supported in part by National Science Foundation grant AST-2108622, NASA Fermi Guest Investigator grants 80NSSC21K1917 and 80NSSC22K1571, and NASA Swift Guest Investigator grant 80NSSC22K1482. This work was supported in part by NSF grant AST-2109127. This study was based

in part on observations conducted using the Perkins Telescope Observatory (PTO) in Arizona, USA, which is owned and operated by Boston University. We acknowledge the use of public data from the Swift data archive. Based on observations obtained with XMM-Newton, an ESA science mission with instruments and contributions directly funded by ESA Member States and NASA. We acknowledge funding to support our NOT observations from the Finnish Centre for Astronomy with ESO (FINCA), University of Turku, Finland (Academy of Finland grant nr 306531). The IAA-CSIC co-authors acknowledge financial support from the Spanish "Ministerio de Ciencia e Innovación" (MCIN/AEI/ 10.13039/501100011033) through the Center of Excellence Severo Ochoa award for the Instituto de Astrofísica de Andalucía-CSIC (CEX2021-001131-S), and through grants PID2019-107847RB-C44 and PID2022-139117NB-C44. The POLAMI observations were carried out at the IRAM 30m Telescope. IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain). Part of the French contribution is supported by the Scientific Research National Center (CNRS) and the French Space Agency (CNES). Some of the data are based on observations collected at the Observatorio de Sierra Nevada, owned and operated by the Instituto de Astrofísica de Andalucía (IAA-CSIC). Further data are based on observations collected at the Centro Astronómico Hispano-Alemán (CAHA), operated jointly by Junta de Andalucía and Consejo Superior de Investigaciones Científicas (IAA-CSIC). CC acknowledges support by the European Research Council (ERC) under the HORIZON ERC Grants 2021 programme under grant agreement No. 101040021. The Submillimetre Array is a joint project between the Smithsonian Astrophysical Observatory and the Academia Sinica Institute of Astronomy and Astrophysics and is funded by the Smithsonian Institution and the Academia Sinica. Mauna Kea, the location of the SMA, is a culturally important site for the indigenous Hawaiian people; we are privileged to study the cosmos from its summit. Some of the data reported here are based on observations made with the Nordic Optical Telescope, owned in collaboration with the University of Turku and Aarhus University, and operated jointly by Aarhus University, the University of Turku, and the University of Oslo, representing Denmark, Finland, and Norway, the University of Iceland and Stockholm University at the Observatorio del Roque de los Muchachos, La Palma, Spain, of the Instituto de Astrofísica de Canarias. E. L. was supported by Academy of Finland projects 317636 and 320045. The data presented here were obtained in part with ALFOSC, which is provided

by the Instituto de Astrofísica de Andalucía (IAA) under a joint agreement with the University of Copenhagen and NOT. S. Kang, S.-S. Lee, W. Y. Cheong, S.-H. Kim, and H.-W. Jeong were supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (MIST) (2020R1A2C2009003). The KVN is a facility operated by the Korea Astronomy and Space Science Institute. The KVN operations are supported by KREONET (Korea Research Environment Open NETwork) which is managed and operated by KISTI (Korea Institute of Science and Technology

Information). Partly based on observations with the 100-m telescope of the MPIfR (Max-Planck-Institut für Radioastronomie) at Effelsberg. Observations with the 100-m radio telescope at Effelsberg have received funding from the European Union’s Horizon 2020 research and innovation programme under grant agreement No 101004719 (ORP).

*Facilities:* Calar Alto, Effelsberg 100-m, Fermi, IRAM-30m, IXPE, KVN, NOT, Perkins, SMA, Swift, T90, XMM-Newton

## REFERENCES

- Abdo, A. A., Ackermann, M., Agudo, I., et al. 2010, *ApJ*, 716, 30, doi: [10.1088/0004-637X/716/1/30](https://doi.org/10.1088/0004-637X/716/1/30)
- Abdollahi, S., Ajello, M., Baldini, L., et al. 2023, *ApJS*, 265, 31, doi: [10.3847/1538-4365/acbb6a](https://doi.org/10.3847/1538-4365/acbb6a)
- Acero, F., Ackermann, M., Ajello, M., et al. 2015, *ApJS*, 218, 23, doi: [10.1088/0067-0049/218/2/23](https://doi.org/10.1088/0067-0049/218/2/23)
- Ackermann, M., Ajello, M., Albert, A., et al. 2015, *ApJL*, 813, L41, doi: [10.1088/2041-8205/813/2/L41](https://doi.org/10.1088/2041-8205/813/2/L41)
- Agudo, I., Thum, C., Ramakrishnan, V., et al. 2018a, *MNRAS*, 473, 1850, doi: [10.1093/mnras/stx2437](https://doi.org/10.1093/mnras/stx2437)
- Agudo, I., Thum, C., Wiesemeyer, H., et al. 2012, *A&A*, 541, A111, doi: [10.1051/0004-6361/201218801](https://doi.org/10.1051/0004-6361/201218801)
- Agudo, I., Marscher, A. P., Jorstad, S. G., et al. 2011a, *ApJL*, 735, L10, doi: [10.1088/2041-8205/735/1/L10](https://doi.org/10.1088/2041-8205/735/1/L10)
- Agudo, I., Jorstad, S. G., Marscher, A. P., et al. 2011b, *ApJL*, 726, L13, doi: [10.1088/2041-8205/726/1/L13](https://doi.org/10.1088/2041-8205/726/1/L13)
- Agudo, I., Thum, C., Molina, S. N., et al. 2018b, *MNRAS*, 474, 1427, doi: [10.1093/mnras/stx2435](https://doi.org/10.1093/mnras/stx2435)
- Ajello, M., Angioni, R., Axelsson, M., et al. 2020, *ApJ*, 892, 105, doi: [10.3847/1538-4357/ab791e](https://doi.org/10.3847/1538-4357/ab791e)
- Aleksić, J., Alvarez, E. A., Antonelli, L. A., et al. 2012, *ApJ*, 748, 46, doi: [10.1088/0004-637X/748/1/46](https://doi.org/10.1088/0004-637X/748/1/46)
- Angel, J. R. P., & Stockman, H. S. 1980, *ARA&A*, 18, 321, doi: [10.1146/annurev.aa.18.090180.001541](https://doi.org/10.1146/annurev.aa.18.090180.001541)
- Angelakis, E., Hovatta, T., Blinov, D., et al. 2016, *MNRAS*, 463, 3365, doi: [10.1093/mnras/stw2217](https://doi.org/10.1093/mnras/stw2217)
- Antonucci, R. 1993, *ARA&A*, 31, 473, doi: [10.1146/annurev.aa.31.090193.002353](https://doi.org/10.1146/annurev.aa.31.090193.002353)
- Arnaud, K. A. 1996, in *Astronomical Society of the Pacific Conference Series*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes, 17
- Aumont, J., Conversi, L., Thum, C., et al. 2010, *A&A*, 514, A70, doi: [10.1051/0004-6361/200913834](https://doi.org/10.1051/0004-6361/200913834)
- Baldini, L., Barbanera, M., Bellazzini, R., et al. 2021, *Astroparticle Physics*, 133, 102628, doi: [10.1016/j.astropartphys.2021.102628](https://doi.org/10.1016/j.astropartphys.2021.102628)
- Baldini, L., Bucciantini, N., Lalla, N. D., et al. 2022, *SoftwareX*, 19, 101194, doi: [10.1016/j.softx.2022.101194](https://doi.org/10.1016/j.softx.2022.101194)
- Bellazzini, R., Angelini, F., Baldini, L., et al. 2003, in *Proc. SPIE*, Vol. 4843, *Polarimetry in Astronomy*, ed. S. Fineschi, 383–393, doi: [10.1117/12.459381](https://doi.org/10.1117/12.459381)
- Blandford, R., Meier, D., & Readhead, A. 2019, *ARA&A*, 57, 467, doi: [10.1146/annurev-astro-081817-051948](https://doi.org/10.1146/annurev-astro-081817-051948)
- Blinov, D., Pavlidou, V., Papadakis, I., et al. 2015, *MNRAS*, 453, 1669, doi: [10.1093/mnras/stv1723](https://doi.org/10.1093/mnras/stv1723)
- Blinov, D., Pavlidou, V., Papadakis, I. E., et al. 2016a, *MNRAS*, 457, 2252, doi: [10.1093/mnras/stw158](https://doi.org/10.1093/mnras/stw158)
- Blinov, D., Pavlidou, V., Papadakis, I., et al. 2016b, *MNRAS*, 462, 1775, doi: [10.1093/mnras/stw1732](https://doi.org/10.1093/mnras/stw1732)
- . 2018, *MNRAS*, 474, 1296, doi: [10.1093/mnras/stx2786](https://doi.org/10.1093/mnras/stx2786)
- Bodo, G., Tavecchio, F., & Sironi, L. 2021, *MNRAS*, 501, 2836, doi: [10.1093/mnras/staa3620](https://doi.org/10.1093/mnras/staa3620)
- Boettcher, M. 2012, *arXiv e-prints*, arXiv:1205.0539, doi: [10.48550/arXiv.1205.0539](https://doi.org/10.48550/arXiv.1205.0539)
- Cerruti, M., Zech, A., Boisson, C., & Inoue, S. 2015, *MNRAS*, 448, 910, doi: [10.1093/mnras/stu2691](https://doi.org/10.1093/mnras/stu2691)
- Costa, E., Soffitta, P., Bellazzini, R., et al. 2001, *Nature*, 411, 662, doi: [10.1038/35079508](https://doi.org/10.1038/35079508)
- Di Gesu, L., Donnarumma, I., Tavecchio, F., et al. 2022, *ApJL*, 938, L7, doi: [10.3847/2041-8213/ac913a](https://doi.org/10.3847/2041-8213/ac913a)
- Di Gesu, L., Marshall, H. L., Ehlert, S. R., et al. 2023, *arXiv e-prints*, arXiv:2305.13497, doi: [10.48550/arXiv.2305.13497](https://doi.org/10.48550/arXiv.2305.13497)
- Di Marco, A., Costa, E., Muleri, F., et al. 2022, *AJ*, 163, 170, doi: [10.3847/1538-3881/ac51c9](https://doi.org/10.3847/1538-3881/ac51c9)
- Di Marco, A., Soffitta, P., Costa, E., et al. 2023, *AJ*, 165, 143, doi: [10.3847/1538-3881/acba0f](https://doi.org/10.3847/1538-3881/acba0f)
- Dorigo Jones, J., Johnson, S. D., Muzahid, S., et al. 2022, *MNRAS*, 509, 4330, doi: [10.1093/mnras/stab3331](https://doi.org/10.1093/mnras/stab3331)
- Ehlert, S. R., Ferrazzoli, R., Marinucci, A., et al. 2022, *ApJ*, 935, 116, doi: [10.3847/1538-4357/ac8056](https://doi.org/10.3847/1538-4357/ac8056)

- Fossati, G., Maraschi, L., Celotti, A., Comastri, A., & Ghisellini, G. 1998, *MNRAS*, 299, 433, doi: [10.1046/j.1365-8711.1998.01828.x](https://doi.org/10.1046/j.1365-8711.1998.01828.x)
- Gabriel, C., Denby, M., Fyfe, D. J., et al. 2004, in *Astronomical Society of the Pacific Conference Series*, Vol. 314, *Astronomical Data Analysis Software and Systems (ADASS) XIII*, ed. F. Ochsenbein, M. G. Allen, & D. Egret, 759
- Gehrels, N., Chincarini, G., Giommi, P., & et al. 2004, *ApJ*, 611, 38, doi: [10.1086/422091](https://doi.org/10.1086/422091)
- Ghisellini, G., Celotti, A., Fossati, G., Maraschi, L., & Comastri, A. 1998a, *MNRAS*, 301, 451, doi: [10.1046/j.1365-8711.1998.02032.x](https://doi.org/10.1046/j.1365-8711.1998.02032.x)
- . 1998b, *MNRAS*, 301, 451, doi: [10.1046/j.1365-8711.1998.02032.x](https://doi.org/10.1046/j.1365-8711.1998.02032.x)
- Gianolli, V. E., Kim, D. E., Bianchi, S., et al. 2023, *MNRAS*, doi: [10.1093/mnras/stad1697](https://doi.org/10.1093/mnras/stad1697)
- Giommi, P., Perri, M., Capalbi, M., et al. 2021, *MNRAS*, 507, 5690, doi: [10.1093/mnras/stab2425](https://doi.org/10.1093/mnras/stab2425)
- HI4PI Collaboration, Ben Bekhti, N., Flöer, L., et al. 2016, *A&A*, 594, A116, doi: [10.1051/0004-6361/201629178](https://doi.org/10.1051/0004-6361/201629178)
- Ho, P. T. P., Moran, J. M., & Lo, K. Y. 2004, *ApJL*, 616, L1, doi: [10.1086/423245](https://doi.org/10.1086/423245)
- Hosking, D. N., & Sironi, L. 2020, *ApJL*, 900, L23, doi: [10.3847/2041-8213/abafa6](https://doi.org/10.3847/2041-8213/abafa6)
- Hovatta, T., & Lindfors, E. 2019, *NewAR*, 87, 101541, doi: [10.1016/j.newar.2020.101541](https://doi.org/10.1016/j.newar.2020.101541)
- Hovatta, T., Valtaoja, E., Tornikoski, M., & Lähteenmäki, A. 2009, *A&A*, 494, 527, doi: [10.1051/0004-6361:200811150](https://doi.org/10.1051/0004-6361:200811150)
- Hovatta, T., Lindfors, E., Blinov, D., et al. 2016, *A&A*, 596, A78, doi: [10.1051/0004-6361/201628974](https://doi.org/10.1051/0004-6361/201628974)
- Ingram, A., Ewing, M., Marinucci, A., et al. 2023, *arXiv e-prints*, arXiv:2305.13028, doi: [10.48550/arXiv.2305.13028](https://doi.org/10.48550/arXiv.2305.13028)
- Jansen, F., Lumb, D., Altieri, B., et al. 2001, *A&A*, 365, L1, doi: [10.1051/0004-6361:20000036](https://doi.org/10.1051/0004-6361:20000036)
- Jethwa, P., Saxton, R., Guainazzi, M., Rodriguez-Pascual, P., & Stuhlinger, M. 2015, *A&A*, 581, A104, doi: [10.1051/0004-6361/201425579](https://doi.org/10.1051/0004-6361/201425579)
- Johnson, S. D., Mulchaey, J. S., Chen, H.-W., et al. 2019, *ApJL*, 884, L31, doi: [10.3847/2041-8213/ab479a](https://doi.org/10.3847/2041-8213/ab479a)
- Kang, S., Lee, S.-S., & Byun, D.-Y. 2015, *Journal of Korean Astronomical Society*, 48, 257, doi: [10.5303/JKAS.2015.48.5.257](https://doi.org/10.5303/JKAS.2015.48.5.257)
- Kraus, A., Krichbaum, T. P., Wegner, R., et al. 2003, *A&A*, 401, 161, doi: [10.1051/0004-6361:20030118](https://doi.org/10.1051/0004-6361:20030118)
- Lähteenmäki, A., & Valtaoja, E. 1999, *ApJ*, 521, 493, doi: [10.1086/307587](https://doi.org/10.1086/307587)
- Lico, R., Liu, J., Giroletti, M., et al. 2020, *A&A*, 634, A87, doi: [10.1051/0004-6361/201936564](https://doi.org/10.1051/0004-6361/201936564)
- Lioudakis, I., Hovatta, T., Huppenkothen, D., et al. 2018a, *ApJ*, 866, 137, doi: [10.3847/1538-4357/aae2b7](https://doi.org/10.3847/1538-4357/aae2b7)
- Lioudakis, I., Romani, R. W., Filippenko, A. V., et al. 2018b, *MNRAS*, 480, 5517, doi: [10.1093/mnras/sty2264](https://doi.org/10.1093/mnras/sty2264)
- Lioudakis, I., Romani, R. W., Filippenko, A. V., Kocevski, D., & Zheng, W. 2019, *ApJ*, 880, 32, doi: [10.3847/1538-4357/ab26b7](https://doi.org/10.3847/1538-4357/ab26b7)
- Lioudakis, I., Blinov, D., Jorstad, S. G., et al. 2020, *ApJ*, 902, 61, doi: [10.3847/1538-4357/abb1b8](https://doi.org/10.3847/1538-4357/abb1b8)
- Lioudakis, I., Marscher, A. P., Agudo, I., et al. 2022, *Nature*, 611, 677, doi: [10.1038/s41586-022-05338-0](https://doi.org/10.1038/s41586-022-05338-0)
- Maraschi, L., Ghisellini, G., & Celotti, A. 1992, *ApJL*, 397, L5, doi: [10.1086/186531](https://doi.org/10.1086/186531)
- Marinucci, A., Muleri, F., Dovciak, M., et al. 2022, *MNRAS*, 516, 5907, doi: [10.1093/mnras/stac2634](https://doi.org/10.1093/mnras/stac2634)
- Marrone, D. P., & Rao, R. 2008, in *Proc. SPIE*, Vol. 7020, *Millimeter and Submillimeter Detectors and Instrumentation for Astronomy IV*, ed. W. D. Duncan, W. S. Holland, S. Withington, & J. Zmuidzinas, 70202B, doi: [10.1117/12.788677](https://doi.org/10.1117/12.788677)
- Marscher, A., Jorstad, S., & Williamson, K. 2017, *Galaxies*, 5, 63, doi: [10.3390/galaxies5040063](https://doi.org/10.3390/galaxies5040063)
- Marscher, A. P., & Gear, W. K. 1985, *ApJ*, 298, 114, doi: [10.1086/163592](https://doi.org/10.1086/163592)
- Marscher, A. P., & Jorstad, S. G. 2021, *Galaxies*, 9, 27, doi: [10.3390/galaxies9020027](https://doi.org/10.3390/galaxies9020027)
- Marscher, A. P., Jorstad, S. G., D’Arcangelo, F. D., et al. 2008, *Nature*, 452, 966, doi: [10.1038/nature06895](https://doi.org/10.1038/nature06895)
- Marscher, A. P., Jorstad, S. G., Larionov, V. M., et al. 2010, *ApJL*, 710, L126, doi: [10.1088/2041-8205/710/2/L126](https://doi.org/10.1088/2041-8205/710/2/L126)
- Maselli, A., Massaro, E., Nesci, R., et al. 2010, *A&A*, 512, A74, doi: [10.1051/0004-6361/200913343](https://doi.org/10.1051/0004-6361/200913343)
- Massaro, E., Perri, M., Giommi, P., & Nesci, R. 2004, *A&A*, 413, 489, doi: [10.1051/0004-6361:20031558](https://doi.org/10.1051/0004-6361:20031558)
- Middei, R., Giommi, P., Perri, M., et al. 2022, *MNRAS*, 514, 3179, doi: [10.1093/mnras/stac1185](https://doi.org/10.1093/mnras/stac1185)
- Middei, R., Lioudakis, I., Perri, M., et al. 2023, *ApJL*, 942, L10, doi: [10.3847/2041-8213/aca281](https://doi.org/10.3847/2041-8213/aca281)
- Myserlis, I., Angelakis, E., Kraus, A., et al. 2018, *A&A*, 609, A68, doi: [10.1051/0004-6361/201630301](https://doi.org/10.1051/0004-6361/201630301)
- Nicastro, F., Kaastra, J., Krongold, Y., et al. 2018, *Nature*, 558, 406, doi: [10.1038/s41586-018-0204-1](https://doi.org/10.1038/s41586-018-0204-1)
- Nilsson, K., Lindfors, E., Takalo, L. O., et al. 2018, *A&A*, 620, A185, doi: [10.1051/0004-6361/201833621](https://doi.org/10.1051/0004-6361/201833621)
- Padovani, P., & Giommi, P. 1995, *ApJ*, 444, 567, doi: [10.1086/175631](https://doi.org/10.1086/175631)
- Padovani, P., Alexander, D. M., Assef, R. J., et al. 2017, *A&A Rv*, 25, 2, doi: [10.1007/s00159-017-0102-9](https://doi.org/10.1007/s00159-017-0102-9)

- Peirson, A. L., Negro, M., Liidakis, I., et al. 2023, *ApJL*, 948, L25, doi: [10.3847/2041-8213/acd242](https://doi.org/10.3847/2041-8213/acd242)
- Perlman, E. S., Biretta, J. A., Zhou, F., Sparks, W. B., & Macchetto, F. D. 1999, *AJ*, 117, 2185, doi: [10.1086/300844](https://doi.org/10.1086/300844)
- Perlman, E. S., & Wilson, A. S. 2005, *ApJ*, 627, 140, doi: [10.1086/430340](https://doi.org/10.1086/430340)
- Pesce-Rollins, M., Lalla, N. D., Omodei, N., & Baldini, L. 2019, *Nuclear Instruments and Methods in Physics Research A*, 936, 224, doi: [10.1016/j.nima.2018.10.041](https://doi.org/10.1016/j.nima.2018.10.041)
- Piconcelli, E., Jimenez-Bailón, E., Guainazzi, M., et al. 2004, *MNRAS*, 351, 161, doi: [10.1111/j.1365-2966.2004.07764.x](https://doi.org/10.1111/j.1365-2966.2004.07764.x)
- Primiani, R. A., Young, K. H., Young, A., et al. 2016, *Journal of Astronomical Instrumentation*, 5, 1641006, doi: [10.1142/S2251171716410063](https://doi.org/10.1142/S2251171716410063)
- Sreekumar, P., Bertsch, D. L., Dingus, B. L., et al. 1998, *ApJ*, 494, 523, doi: [10.1086/305222](https://doi.org/10.1086/305222)
- Strüder, L., Briel, U., Dennerl, K., et al. 2001, *A&A*, 365, L18, doi: [10.1051/0004-6361:20000066](https://doi.org/10.1051/0004-6361:20000066)
- Tagliacozzo, D., Marinucci, A., Ursini, F., et al. 2023, arXiv e-prints, arXiv:2305.10213, doi: [10.48550/arXiv.2305.10213](https://doi.org/10.48550/arXiv.2305.10213)
- Tavani, M., Cavaliere, A., Munar-Adrover, P., & Argan, A. 2018, *ApJ*, 854, 11, doi: [10.3847/1538-4357/aaa3f4](https://doi.org/10.3847/1538-4357/aaa3f4)
- Tavecchio, F., Landoni, M., Sironi, L., & Coppi, P. 2018, *MNRAS*, 480, 2872, doi: [10.1093/mnras/sty1491](https://doi.org/10.1093/mnras/sty1491)
- Thum, C., Agudo, I., Molina, S. N., et al. 2018, *MNRAS*, 473, 2506, doi: [10.1093/mnras/stx2436](https://doi.org/10.1093/mnras/stx2436)
- Urry, C. M., & Padovani, P. 1995, *PASP*, 107, 803, doi: [10.1086/133630](https://doi.org/10.1086/133630)
- Ursini, F., Marinucci, A., Matt, G., et al. 2023, *MNRAS*, 519, 50, doi: [10.1093/mnras/stac3189](https://doi.org/10.1093/mnras/stac3189)
- Weisskopf, M. C., Soffitta, P., Baldini, L., et al. 2022, *Journal of Astronomical Telescopes, Instruments, and Systems*, 8, 026002, doi: [10.1117/1.JATIS.8.2.026002](https://doi.org/10.1117/1.JATIS.8.2.026002)
- Zhang, H., Marscher, A. P., Guo, F., et al. 2023, *ApJ*, 949, 71, doi: [10.3847/1538-4357/acc657](https://doi.org/10.3847/1538-4357/acc657)